



DEPDC5 gene

DEP domain containing 5

Normal Function

The *DEPDC5* gene provides instructions for making a protein that is one piece of a group of proteins (complex) called GATOR1. This complex is found in cells throughout the body, where it regulates a signaling pathway called the mTOR pathway. The mTOR pathway is involved in cell growth and division (proliferation), the survival of cells, and the creation (synthesis) of new proteins. The role of the GATOR1 complex is to block this pathway by inhibiting (stopping) the activity of a complex called mTOR complex 1 (mTORC1) that is integral to the mTOR pathway.

In the brain, the mTOR pathway regulates many processes, including the growth and development of nerve cells and their ability to change and adapt over time (plasticity).

Health Conditions Related to Genetic Changes

familial focal epilepsy with variable foci

More than 80 mutations in the *DEPDC5* gene have been found to cause familial focal epilepsy with variable foci (FFEVF), which is an uncommon form of recurrent seizures (epilepsy) that runs in families. Affected individuals experience focal seizures, which are seizures that do not cause a loss of consciousness. Most of the *DEPDC5* gene mutations lead to the production of an abnormally short protein that is quickly broken down. As a result, formation of normal GATOR1 complex is reduced, leading to overactivity of mTORC1 and excessive signaling of the mTOR pathway. It is not clear how an abnormally active mTOR pathway leads to the focal seizures of FFEVF. Research suggests that increased mTOR pathway signaling in the brain leads to changes in the connections between nerve cells (synapses) and increased activation (excitation) of nerve cells, which can cause seizures.

For unknown reasons, some people with FFEVF caused by a *DEPDC5* gene mutation never develop the condition, a situation known as reduced penetrance. It is estimated that 60 percent of individuals with *DEPDC5* gene mutations go on to develop FFEVF.

other disorders

Mutations in the *DEPDC5* gene can cause other seizure disorders, known as familial mesial temporal lobe epilepsy and infantile spasms. Similar to individuals with FFEVF (described above), people with familial mesial temporal lobe epilepsy have focal seizures. They may also have feelings of déjà vu, fear, or nausea during the seizure.


Infantile spasms are seizures that usually appear before the age of 1 and involve recurrent muscle contractions.

As in FFEVF, most of the *DEPDC5* gene mutations that cause familial mesial temporal lobe epilepsy or infantile spasms lead to reduced GATOR1 complex formation and an abnormally active mTOR pathway. It is unclear why individuals with mutations in the same gene develop different seizure disorders.

Chromosomal Location

Cytogenetic Location: 22q12.2-q12.3, which is the long (q) arm of chromosome 22 between positions 12.2 and 12.3

Molecular Location: base pairs 31,753,951 to 31,907,034 on chromosome 22 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- DEP.5
- FFEVF
- FFEVF1
- KIAA0645

Additional Information & Resources

Educational Resources

- Jasper's Basic Mechanisms of the Epilepsies (fourth edition, 2012): mTOR and Epileptogenesis in Developmental Brain Malformations
<https://www.ncbi.nlm.nih.gov/books/NBK98145/>

GeneReviews

- DEPDC5-Related Epilepsy
<https://www.ncbi.nlm.nih.gov/books/NBK385626>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28DEPDC5%5BTIAB%5D%29+OR+%28DEP+domain+containing+5%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- DEP DOMAIN-CONTAINING PROTEIN 5
<http://omim.org/entry/614191>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_DEPDC5.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=DEPDC5%5Bgene%5D>
- HGNC Gene Family: GATOR1 subcomplex
<http://www.genenames.org/cgi-bin/genefamilies/set/1394>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=18423
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/9681>
- UniProt
<http://www.uniprot.org/uniprot/O75140>

Sources for This Summary

- Baldassari S, Licchetta L, Tinuper P, Bisulli F, Pippucci T. GATOR1 complex: the common genetic actor in focal epilepsies. *J Med Genet.* 2016 Aug;53(8):503-10. doi: 10.1136/jmedgenet-2016-103883. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/27208208>
- Baulac S, Ishida S, Marsan E, Miquel C, Biraben A, Nguyen DK, Nordli D, Cossette P, Nguyen S, Lambrecq V, Vlaicu M, Daniau M, Bielle F, Andermann E, Andermann F, Leguern E, Chassoux F, Picard F. Familial focal epilepsy with focal cortical dysplasia due to DEPDC5 mutations. *Ann Neurol.* 2015 Apr;77(4):675-83. doi: 10.1002/ana.24368. Epub 2015 Mar 13.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25623524>
- Baulac S. Genetics advances in autosomal dominant focal epilepsies: focus on DEPDC5. *Prog Brain Res.* 2014;213:123-39. doi: 10.1016/B978-0-444-63326-2.00007-7. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25194487>

- Baulac S. mTOR signaling pathway genes in focal epilepsies. *Prog Brain Res.* 2016;226:61-79. doi: 10.1016/bs.pbr.2016.04.013.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/27323939>
- OMIM: DEP DOMAIN-CONTAINING PROTEIN 5
<http://omim.org/entry/614191>
- Dibbens LM, de Vries B, Donatello S, Heron SE, Hodgson BL, Chintawar S, Crompton DE, Hughes JN, Bellows ST, Klein KM, Callenbach PM, Corbett MA, Gardner AE, Kivity S, Iona X, Regan BM, Weller CM, Crimmins D, O'Brien TJ, Guerrero-López R, Mulley JC, Dubeau F, Licchetta L, Bisulli F, Cossette P, Thomas PQ, Gecz J, Serratosa J, Brouwer OF, Andermann F, Andermann E, van den Maagdenberg AM, Pandolfo M, Berkovic SF, Scheffer IE. Mutations in DEPDC5 cause familial focal epilepsy with variable foci. *Nat Genet.* 2013 May;45(5):546-51. doi: 10.1038/ng.2599.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23542697>
- Ishida S, Picard F, Rudolf G, Noé E, Achaz G, Thomas P, Genton P, Mundwiller E, Wolff M, Marescaux C, Miles R, Baulac M, Hirsch E, Leguern E, Baulac S. Mutations of DEPDC5 cause autosomal dominant focal epilepsies. *Nat Genet.* 2013 May;45(5):552-5. doi: 10.1038/ng.2601. Epub 2013 Mar 31.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23542701>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5010101/>
- Scheffer IE, Heron SE, Regan BM, Mandelstam S, Crompton DE, Hodgson BL, Licchetta L, Provini F, Bisulli F, Vadlamudi L, Gecz J, Connelly A, Tinuper P, Ricos MG, Berkovic SF, Dibbens LM. Mutations in mammalian target of rapamycin regulator DEPDC5 cause focal epilepsy with brain malformations. *Ann Neurol.* 2014 May;75(5):782-7. doi: 10.1002/ana.24126.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24585383>
- Weckhuysen S, Marsan E, Lambrecq V, Marchal C, Morin-Brureau M, An-Gourfinkel I, Baulac M, Fohlen M, Kallay Zetchi C, Seeck M, de la Grange P, Dermaut B, Meurs A, Thomas P, Chassoux F, Leguern E, Picard F, Baulac S. Involvement of GATOR complex genes in familial focal epilepsies and focal cortical dysplasia. *Epilepsia.* 2016 Jun;57(6):994-1003. doi: 10.1111/epi.13391.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/27173016>
- van Kranenburg M, Hoogeveen-Westerveld M, Nellist M. Preliminary functional assessment and classification of DEPDC5 variants associated with focal epilepsy. *Hum Mutat.* 2015 Feb;36(2):200-9. doi: 10.1002/humu.22723.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25366275>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/DEPDC5>

Reviewed: March 2017

Published: April 25, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services